

PHARMACY ORDERS
Anorectal Surgery

Name: _____ DOB: _____ Date: _____

1. Donut pillow for sitting in a chair.
2. Non-sterile latex medical gloves, small package.
3. Thick quilted baby wipes
4. Analpram® H.C. 2.5% 1 oz., 2 refills
Apply locally t.i.d.
5. 4" x 4" non-sterile gauze pads, small package.
6. Maxi Pads, 1 package
7. Demerol® 100mg. ampules, Disp: #2 ampules
Demerol® 50-100mg + Vistaril® 25-50mg q3-4 h prn pain. If analgesia is relatively unsatisfactory, call physician.
8. Vistaril® 50mg./cc, Disp: #1 small vial
Demerol® 50-100mg + Vistaril® 25-50mg q3-4 h prn pain. If analgesia is relatively unsatisfactory, call physician.
9. In the event that a CADD-PCA pump is prescribed, do not dispense Demerol® and Vistaril®, but instead dispense:

Morphine Sulfate 25mg/cc, 20ml vial, Disp: #5
Morphine Sulfate, subcutaneous infusion by CADD-PCA pump, 0.8 mg/hr basal flow rate with bolus dose 0.2 mg q h, as indicated by patient response. The maximum number of dosage increases of 0.2mg each that is allowed before calling the pharmacist is 3. The pharmacist has authority to increase the dosage if analgesia is inadequate.
10. Dilaudid® tablets, Disp: #100
1 tabs q4 h prn pain. 1 additional tab q45-60 min. prn breakthrough pain.
11. Syringes #2, 23 gauge 1 inch
12. Alcohol wipes #5
13. Pericolace®, Disp: #20
2 hs. prn BM.

14. Ambien® 10mg, Disp: #10, 1 refill
10mg hs prn sleep. May repeat once after 2h.
15. Fleets® Saline Sodium Phosphate Enema, Disp: 2
16. Fleets® Mineral Oil Enema, Disp: 1
17. Cotton roll, Disp: 1
18. Pyridium® 200mg. capsules for male patients only. Disp: #6, 1 t.i.d.
19. Tylenol® 10gr., Disp: small bottle
Counsel patient not to take NSAIDs 7 days before or after surgery.

PHYSICIAN SIGNATURE